



# IgG4ward!

PAVING A PATH 4WARD

## Vaccine Guidelines

### General Vaccination Recommendations:

- **Optimize Timing:**
  - When possible, schedule vaccinations during periods of lower immunosuppression to ensure a stronger immune response.
- **B-cell Depleting Therapy:**
  - Plan vaccinations before scheduled B-cell depletion therapy. Avoid vaccinating immediately after receiving therapy. (B-cells are important to achieving normal responses to vaccines. The response to the vaccine may be muted if you have just been treated with a B-cell depleting medication).
  - If receiving intermittent immunosuppression (for example, B-cell depletion every once in a while), aim to be vaccinated 2 to 4 weeks before the first infusion.
- **Partial Protection is Better than None:**
  - If unvaccinated, it is still advisable to receive vaccines even during immunosuppression, knowing revaccination (a booster) may be necessary later.

*Disclaimer:*

*This document provides general vaccine guidance for immunocompromised individuals, specifically those on B-cell depleting therapies. It is not intended to replace professional medical advice. Individual vaccine recommendations should be discussed with a healthcare provider based on personal health*



# IgG4ward!

PAVING A PATH 4WARD

- **Booster Doses:**

- Consider booster doses or revaccination as needed.
  - **Example:** COVID-19 vaccines may be recommended every 6 months rather than yearly for immunocompromised individuals.

- **Multiple Vaccines in One Visit:**

- It is safe to receive more than one vaccine on the same day, but mild side effects like fatigue or arm pain may occur.
- To manage vaccine site soreness: Use acetaminophen (Tylenol) or ibuprofen (Motrin/Advil) – if this is approved by your clinical team. Can also use an ice pack.

## Key Vaccine Guidelines:

- Influenza (Flu) Vaccine:

- Inactivated (IIV) or Recombinant (RIV) only
- **Timing:**
  - At least 4 weeks before starting B-cell depleting therapy for optimal response.
  - If already on therapy, vaccinate 5-6 months after the last infusion and at least 4 weeks before the next infusion, if possible.
  - **Avoid:** Live nasal spray vaccine (LAIV)

*Disclaimer:*

*This document provides general vaccine guidance for immunocompromised individuals, specifically those on B-cell depleting therapies. It is not intended to replace professional medical advice. Individual vaccine recommendations should be discussed with a healthcare provider based on personal health*



# IgG4ward!

PAVING A PATH 4WARD

- **COVID-19 Vaccine:**

- **Timing:**

- If already on B-cell depletion therapy, administer 2-4 weeks before the next scheduled infusion.
- Revaccination may be needed 6 months after therapy completion if vaccinated while immunosuppressed.

- **Boosters:**

- May be needed 6 months instead of annually—discuss with your healthcare team.

- **Prednisone Considerations:**

- If on long-term prednisone, vaccination is ideal at 5-10mg daily, but if higher doses are required, vaccination is still recommended with a discussion with your healthcare team about booster strategies.

- **Respiratory Syncytial Virus (RSV) Vaccine:**

- **Recommended for:**

- All adults aged 75+
- Adults aged 60-74 with increased RSV risk (including moderate/severe immune compromise)
- Not yet recommended for those under 60 years old

**Disclaimer:**

This document provides general vaccine guidance for immunocompromised individuals, specifically those on B-cell depleting therapies. It is not intended to replace professional medical advice. Individual vaccine recommendations should be discussed with a healthcare provider based on personal health



# IgG4ward!

PAVING A PATH 4WARD

- Pneumococcal Vaccine (Pneumonia)

- **Recommended for:**

- Adults  $\geq 19$  years with immunosuppression

- **Timing:**

- Preferably before therapy; if given during therapy, immune response may be lower

- Shingles (Herpes Zoster) Vaccine

- **Risk:** Immunocompromised individuals are 51% more likely to develop shingles

- **Recommended for:**

- All immunocompromised adults  $\geq 18$  years old (CDC-approved, even if under 50—pharmacists may require CDC documentation)

- **Timing:**

- Two doses 2-6 months apart (can be longer if needed to avoid intense immunosuppression periods)

**Disclaimer:**

This document provides general vaccine guidance for immunocompromised individuals, specifically those on B-cell depleting therapies. It is not intended to replace professional medical advice. Individual vaccine recommendations should be discussed with a healthcare provider based on personal health



# IgG4ward!

PAVING A PATH 4WARD

## ● Additional Considerations

- Consult your healthcare team to discuss individual vaccine timing based on your disease activity and treatment schedule.
- Vaccination has been shown to reduce risk of severe disease and complications even if the immune response is lower.
- Revaccination should be considered for individuals who received vaccines while on B-cell depleting therapy.

### Disclaimer:

This document provides general vaccine guidance for immunocompromised individuals, specifically those on B-cell depleting therapies. It is not intended to replace professional medical advice. Individual vaccine recommendations should be discussed with a healthcare provider based on personal health